

THIR UNITED STRATES OF AMIERICA

<u>co am ro whom rhese; presents; sham come;</u> Agriculture and Agri-Jood Canada and Misconsin Alumni Research Joundation

MOCCUS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HERS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY EARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC LENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR UNG IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE REPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT ATHE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

POTATO

'AC Glacier Chip'

In Jestimonn Thereof. I have hereunto set my hand and caused the seal of the Plant Hariety Frotestion Office to be affixed at the City of Washington, D.C. this seventh day of February, in the year two thousand and eight.

Atlest:

JCmZ~

Commissioner Plant Variety Protection Office Agricultural Marketing Service Colmand T: Schaff

Director

S&T-470 (2-99) designed by the Plant Variety Protection Office with WordPerfect 6.0a. Replaces STD-470 (6-98) which is obsolete.

Form Approved - OMB No. 0581-0055

U.S. DEPARTMENT OF AGRICULTURE
AGRICULTURAL MARKETING SERVICE
SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE

The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995.

Application is required in order to determine if a plant veriety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).

(See reverse for instructions and information coll

APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE (Instructions and information collection burden statement on reverse) 2. TEMPORARY DESIGNATION OR EXPERIMENTAL NAME NAME OF OWNER Agriculture and Agri-Food Canada and Wisconsin 3. VARIETY NAME COST espondance - NAC Alumni Research Foundation WIS 75-30 AC Glacier Chip FOR OFFICIAL USE ONLY 4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Country) 5. TELEPHONE (include area code) Director, Agriculture and Agri-Food Canada, 5403-1st **PVPO NUMBER** Ave South, Lethbridge, Alberta, Canada T1J 4B1 403-327-4561 aoo100043 6. FAX (include area code) FILING DATE 403-382-3156 9. DATE OF INCORPORATION IF THE OWNER NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZATION (corporation, partnership, association, etc.) 8. IF INCORPORATED, GIVE STATE OF INCORPORATION 12/29/00 Dept. of Canadian Federal Govt. N/A N/A FILING AND EXAMINATION FEES: 10. NAME AND ADDRESS OF OWNER REPRESENTATIVE(S) TO SERVE IN THIS APPLICATION. (First person listed will receive all papers) c/o Dr. S.D. Morgan Jones Director Agriculture and Agri-Food Canada Research Centre 5403-1st Ave South Lethbridge, Alberta Canada T1J 4B1 11. TELEPHONE (Include area code) 12. FAX (Include area code) 13 F-MAII 14. CROP KIND (Common Name) Potato 403-327-4561 403-382-3156 morganjoness@em.agr.ca 19. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE SOLD AS A CLASS OF 18. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow instructions on See Section 83(a) of the Plant Variety Protection Act) CERTIFIED SEED? reverse) NO (if "no," go to item 22) X. Exhibit A. Origin and Breeding History of the Variety YES (If "yes", enswer items 20 and 21 below) Exhibit B. Statement of Distinctness 20. DOES THE OWNER SPECIFY THAT SEED OF THIS YES VARIETY BE LIMITED AS TO NUMBER OF CLASSES? Exhibit C. Objective Description of Variety IF YES, WHICH CLASSES? FOUNDATION REGISTERED CERTIFIED Exhibit D. Additional Description of the Variety (Optional) Exhibit E. Statement of the Basis of the Owner's Ownership 21. DOES THE OWNER SPECIFY THAT THE CLASSES BE Voucher Sample (2,500 viable untreated seeds or, for tuber propagated varieties, LIMITED AS TO NUMBER OF GENERATIONS? NO verification that tissue culture will be depositied and maintained in an approved public IF YES, SPECIFY THE Filing and Examination Fee (\$2,450), made payable to "Treasurer of the United States" (Mail to the Plant Variety Protection Office) NUMBER 1, 2, 3, etc. FOUNDATION REGISTERED CERTIFIED See explanation on next page 23. IS THE VARIETY OR ANY COMPONENT OF THE VARIETY PROTECTED BY INTELLECTUAL 22. HAS THE VARIETY (INCLUDING ANY HARVESTED MATERIAL) OR A HYBRID PRODUCED PROPERTY RIGHT (PLANT BREEDER'S RIGHT OR PATENT)? FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRANSFERRED, OR USED IN THE U. S. OR OTHER COUNTRIES? X YES X IF YES, GIVE COUNTRY, DATE OF FILING OR ISSUANCE AND ASSIGNED IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE, DISPOSITION, TRANSFER, OR USE FOR EACH COUNTRY AND THE CIRCUMSTANCES. (Please use space indicated on reverse.) REFERENCE NUMBER. (Please use space indicated on reverse.) 24. The owners declare that a viable sample of basic seed of the variety will be furnished with application and will be replenished upon request in accordance with such regulations as may be applicable, or for a tuber propagated variety a tissue culture will be deposited in a public repository and maintained for the duration of the certificate. The undersigned owner(s) is(are) the owner of this sexually reproduced or tuber propagated plant variety, and believe(s) that the variety is new, distinct, uniform, and stable as required in Section 42, and is entitled to protection under the provisions of Section 42 of the Plant Variety Protection Act. Owner(s) is(are) informed that false representation herein can jeopardize protection and result in penalties. SIGNATURE OF OWNER SIGNATURE OF OWNER NAME (Please print or type) Dr. S.D. Morgan Jones BRYAN Z. RENK CAPACITY OR TITLE DATE

INSTRUCTIONS

GENERAL: To be effectively filed with the Piant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$2,450 (\$300 filing fee and \$2,150 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 500, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$300 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvp.htm

ITEM

18a. Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 21. See Section 83 of the Act for the Contents and Term of Plant Variety Protection.
- 22. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 5.5 of the Act for instructions on claiming the benefit of an earlier filing date.
- 21. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)

Limited to six field generations beyond stocks laboratory tested to confirm pathogen free status which may be tissue culture plantlets or greenhouse produced minitubers.

22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

AC Glacier Chip was sold to Options Inc. (Connell, WA) on May 31, 2000 (44,880 CWT) for a commercial scale trial.

23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

Accepted for filing in Canada on June 25, 2000 with application #99-1618 (July 2000 issue of The Plant Varieties Journal page 15 of The Canadian Plant Breeders Rights Office).

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the variety names proposed by contacting: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center-East, Beltsville, MD 20705. Telephone: (301) 504-8089.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this collection of information is (0581-0055). The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

DRAFT Exhibit A Form

1. Describe the genealogy (back to and including public and commercial varieties, lines, or clones used) and the breeding method(s).
the cross [wischip (wis 55-232.58x Wis 231) X Wis 69-49.74 (WIS 63-508.65x
Level 7 ms souds by the University of Wiscowsia 4, 1025 and received
by the MFC lethbridge Research Centre in 1980 after 3 years of field
a lastic a Wheen in the close was single and the section of
selection in Wiscousin. the clone was riveleded in the regional
trial system of the Lethbuidge Research Centre breeding program
from 1980-1988. Selection criteria included tuber field, chipping
2. Give the details of subsequent stages of selection and multiplication.
Year Detail of Stage Selection Criteria
1939 Industry evaluation in Alberta, Chipping quality
1997 Large scale Trial by the Potats Grovers of Alberta
Grover of seperta
7,,000
3a. Is the variety uniform? Vyes No
How did you test for uniformity?
How did you test for uniformity? Morphological uniformity was assessed in field trials (Regional Trials) conducted from 1980-1988.
(Regional Trials) conducted from 1980 -1988.
3b. Is the variety stable? Yes No
The presence of variouts was assessed in the field on from tissue culture stocks since testing commenced at the letheridge Research
culture stocks since testing commenced set the appropriate
centre in 1980. No variouts have been identified.
4. Are genetic variants observed or expected during reproduction and multiplication? Yes No
If yes, state how these variants may be identified, their type and frequency.

Exhibit A

Origin and Breeding History of the Variety

AC Glacier Chip was registered in Canada by the Variety Registration Office of the Canadian Food Inspection Agency on June 12, 2000 (Certificate #5149). The original cross was made in 1975 by the University of Wisconsin (Wischip (female) and WIS69-49.74 (male)). AC Glacier Chip was included in a breeding population of clones provided to Agriculture and Agri-Food Canada's Lethbridge Research Centre (AAFC-LRC) by the University of Wisconsin Potato Breeding Program in 1980. The final selection was made by AAFC-LRC.

The variety was included in the regional trials of the Western Canadian Potato Breeding Program for 8 years under the seedling designation WIS75-30. These replicated trials are conducted at trial sites in Alberta, Saskatchewan and Manitoba. Evaluation (research plots) by the Western Canadian Potato Industry commenced in 1989. In 1996 the variety (WIS75-30) was offered to the Western Potato Consortium (Industry/AAFC-LRC partnership for market development of Lethbridge Research Centre potato varieties) and the exclusive right to commercialize was assigned to the Potato Growers of Alberta (6008-46th Avenue, Taber, Alberta, Canada T1G 2B1).

In 1997, 1998 and 1999 AC Glacier Chip was included in replicated multi-harvest trials in Alberta by AAFC-LRC and chip quality from the final harvest stage monitored out of storage for up to 8 months after the final harvest. In 1998 and 1999 the variety was included in the North Central Trials (NCR-84) at several sites in the United States of America and Canada. Pathogen free tissue cultures available from Agriculture and Agri-Food Canada (Lethbridge Research Centre), have been provided to the Potato Growers of Alberta and used to produce an Elite seed source in Alberta, Canada. Quantitative data from the Western Canadian Regional Trials, North Central Trial and the multi-harvest trials are not presented as a part of this application. In 2000 a large scale trial of AC Glacier Chip was conducted in the USA by the Potato Growers of Alberta.

AC Glacier Chip under the denomination WIS75-30 was accepted for filing in Canada on the 25th of June, 1999 (Application # 99-1681 in the July 1999 issue of the Plant Varieties Journal page 15). The denomination has since been changed to AC Glacier Chip, field inspection completed and the detailed description published in the July, 2000 issue (page 41) of the Plant Varieties Journal. The Plant Breeder's Right will be jointly held by Agriculture and Agri-Food Canada and the University of Wisconsin.

No variants have been identified either in the field or from tissue-cultured stocks since the original selection in 1989 indicating that AC Glacier Chip is uniform and stable.

Dr. S.D. Morgan Jones, Director

DRAFT Exhibit B Form


Based on overall morphology, W15 75-30 is a	nost similar to SNOW DEN
Applicant's new variety	Most similar comparison variety(ies)
••••	
WIS 75-30 most clearly differs from SNOW	in the following traits:
Applicant's new variety Most simi	lar comparison variety(ies)

Name the specific trait, then list the value of that trait for each variety in the comparison. Attach appropriate supporting evidence (see the Guidelines for Presenting Evidence in Support of

Variety Distinctness, av	vailable from the PVP Office	or website).	
Eg. Terminal leaflet tip shape Eg. Corolla inner Color	Cuspitate Violet (85A)	Obtuse Red Purple (74B)	photograph attached Royal Horticultural Society Colour Chart
Eg. NumberEye/Tuber	15 +/- 2 (N=100)	30 +/- 4 (N=100)	statistics attached
1. Qualitative traits:	Applicant's New Variety W15 75-30	I* Comparison Variety Snowden	Location of Evidence
- Growth Habit	speading	Semi-erect	Photo attached
-Tuber Shape	oval	Round	Photo Attached
_Light Strout Shape	Conical	ovoid	Photo Attached
2. Color traits:	0-1504	60000	Photo attached
Light grout Base	Ked-Viblet	Green	
_ Light spart Tip	Red-Violet	Green	Photo attached
·			
3. Quantitative traits:			
4. Other:	0		DO + Hooked
Tuber Skin Texture	_ Smooth	Flaky	Photo attached
		weak	Photo attached
_ Lipert Sport Box prebescence	medium	W 22 -	,

Snowden - Tuber String Texture


#200100043



SNOWDEN



#200100043




SNOWDEN






Snowden- crowth Hobet

ORSISIA LIED VIEW ON 1979 DA





FORM APPROVED - OMB NO. 0581-0055

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY DIVISION PLANT VARIETY PROTECTION OFFICE

EXHIBIT C OBJECTIVE DESCRIPTION OF VARIETY POTATO (Solanum tuberosum L.)

Public reporting burden for this collection of information is estimated to average __ minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the form. Send comments regarding this burden estimate or any other aspects of this collection of information, including suggestions for reducing this burden, to USDA, OIRM, Clearance Officer, AG Box 7630, Washington, DC 20250, regarding OMB No. 0581-0055. When replying, refer to OMB number and form number you your letter.

INSTRUCTIONS

The Objective Description Form:

The objective description form lists characteristics to be used as the basis for developing the description of potato varieties. It is designed to guide the applicant in describing a variety in detail so a meaningful comparison with other potato varieties can be accomplished. It is recommended that this form be completed in as much detail as possible to ensure an accurate description. Please fill in the requested data and place the appropriate number that describes the varietal characters typical of this potato variety and the reference varieties in the respective boxes.

Test Guidelines:

Any statistical and trial (field test) data that may be necessary to support the variety description should be attached to this form. Please include for trial data the plot size, number of replications, number of plants, plant spacing, trial locations and growing periods. Trials should normally be conducted at one place, in the region that the variety has been adapted for, with a minimum of one growing period in the United States. All comparative data should be determined from varieties entered in the same trials. The size of the plots should be such that plants or parts of plants may be removed for measuring and counting without prejudice to the observations which must be made at the end of the growing period. As a minimum, each test should include a total of 60 plants which should be divided between two or more replicates. Separate plots for observation and measuring can only be used if they have been subject to similar environmental conditions. To determine color for a plant or plant parts a recognized standard color chart must be used such as the Royal Horticultural Society (R.H.S.) Color Chart.

Reference Varieties:

The application variety should be compared to at least one reference variety preferably a set of reference varieties. The reference varieties should be market class standard varieties currently grown in the United States and or the variety(ies) most similar. The following varieties are recommended as market class standards to be used as reference varieties:

Yellow-flesh table-stock	Yukon Gold
Round-white table-stock	Superior
Chip-processing	Atlantic, Snowden, Norchip
Frozen-processing	Russet Burbank
Russet table-stock	Russet Burbank, Russet Norkotah, Goldrush
Red table-stock	Red Pontiac, Red Norland, Red Lasoda

If the applicant does not use one of the recommended reference varieties the PVP office may not have a complete description for the reference variety used; therefore the applicant may have to supply this description by completing an Exhibit C form for the reference variety.

Characteristics:

The plant type and growth habit characteristics are collected at early first bloom. Figure 1 is supplied to help visualize the growth habit. For this descriptor, look at the stems rather than the stems and foliage. Plant maturity is measured at natural vine senescence.

Stem characteristics are also collected at early bloom. Stem anthocyanin coloration is divided into two descriptors: Location and intensity. Figure 12 is supplied to give an example of stem wings.

STD-476 (01-96)

Page 1 of 19

Leaf characteristics are observed at early first bloom. Fully-developed leaves located on the middle third of the plant should be used. Leaf pubescence refers to general trichomes. Figure 2 is supplied for examples of leaf silhouette. Figure 3 should be used to describe terminal and primary leaflet shape. Figures 4 and 5 are used to describe the terminal and primary leaflet shape of tip and base, respectively. To measure the total number of primary leaflets pairs, collect 10 fully- developed petioles (with leaves attached from each replication and take the average number of secondary and tertiary leaflets. Figure 11 is supplied to define leaf characteristics. Glandular trichomes should be described through descriptor #12 (Additional Comments and Characteristics). Leaf stipules are shown in figure 13 for visual definition.

Inflorescence characteristics should be measured at early first bloom. Figures 6 and 7 are supplied to describe corolla and anther shape, respectively. Corolla, calyx, anther, stigma and pollen should be observed on newly opened flowers.

Berry production should be based on field-grown plants rather than greenhouse plants.

Tuber characteristics should be observed following harvest. Figures 9 and 10 are available to describe distribution of secondary color and tuber shape, respectively.

Disease and pest reactions should be based upon specific tests rather than field observations. Other diseases or pests reactions not requested can be described if it is felt that it would be helpful to the description.

Quality characteristics should be described according to the market use.

If the plant is transgenic, this gene insertion(s) should be described.

Chemical identification and any other characteristics can be describe if they are helpful in distinguishing the variety.

A rating system of 1-9 provides a scale for describing most characteristics in this form. Characteristic may be rated with intermediate values where the characteristic grades gradually from one extreme to another. For example, if the character states are described as: 3 = Small; 5 = Medium; 7 = Large; the other values of 1, 2, 4, 6, 8, or 9 may be selected.

Legend:

V = Application Variety

R1-R4 = Reference Varieties

* = Both the reference variety(ies) and application variety must be described for characteristics designated with an asterisk.

NAME OF APPLICANT(S)		6 (0 2 B)	-07 F	OR OFFICIAL USE ONLY
Agriculture and Agri-Fo Alumni Research Four	4	NUMBER		
ADDRESS (Street and No. or R.F.D. No., City, State, and Zip Code)				00/00043
THE DIRECTOR	.F.D. No., Cuy, State, ana Zij	Code)		ETY (V) NAME
AGRICULTURE A	ND AGRI-FOOD CA	NA DA	AL	GLACIER CHIP
research cen Lethbridge,	TTRE, 1.0. BOX 3000)	1	ORARY OR EXPERIMENTAL NATION
CAMADA TI	184 7			w1575-30
REFERENCE VAR	IETIES: Enter the r	eference vario	ety name in	the appropriate box
Reference Variety 1 (R1)	Reference Variety 2 (R2)	Reference V	ariety 3 (R3)	Reference Variety 4 (R4)
SHOWDEH				
1. MARKET CHARACTERIS	TICS:			
MARKET CLASS:				
1 = Yellow-flesh tables5 = Russet tablestock;	tock; 2 = Round-white table 6 = Other	stock; 3 = Chip-p	rocessing; 4 = 1	Frozen-processing;
v 3	R1 3 R		R3	R4
2. PLANT CHARACTERISTI	CS:			
GROWTH HABIT: (See 3 = Erect (>45° with	figure 1) ground); 5 = Semi-erect (30	-45° with ground);	7 = Spreading.	
v 7	R1 5 R2		R3	R4
TYPE:	n, stems clearly visible); 2 =	Intermediate: 3 —	Leaf (Foliage of	osed stoms handly visible)
1 — Steat (toninge oper	, stems cicarly visible), 2 =	mermenate, 3 –	Lear (Poliage Ci	oseu, stems naruly visione)
v 1	R1 3 R2		R3	R4
MATURITY: Days after	planting (DAP) at vine seneso	cence		,
v il8	R1 118 R2		R3	R4
PLANTING DATE:				
V MAY 14, 1999 R1 5	(291. 8, 1999 R2	R	3	R4
REGION/AREA:				
V VAUXHALL , RI AI	AUXHALL, BERTA, CANADA	R	3	R4
		ـ ـ		

MATURITY CLASS: 1 = Very Early (<1) 5 = Very Late (>1)	100 DAP); 2 = Early (10	0-110 DAP); 3 = Mid-s	eason (111-120 DAP); 4 :	= Late (121-130 DAP);
v 3	R1 3	R2	R3	R4
. STEM CHARACTERIS	IICS: Measure at early fi	rst bloom		
*			· =	<u></u>
STEM ANTHOCYANI 1 = Absent; 3 = W	N COLORATION: /eak; 5 = Medium; 7 =	Strong: 9 = Very Stro	ne	
· · · · · · · · · · · · · · · · · · ·		[] [] [] [] [] [] [] [] [] []	[]	
V	R1	R2	R3	R4
· · · · · · · · · · · · · · · · · · ·	eak; 5 = Medium; 7 =			
v 5	R1 /	R2	R3	R4
LEAF CHARACTERIST	CS:			
1 = Yellowish-green		Medium green; 4 = Da	rk green; 5 = Grey-gree	n; 6 = Other
v 3	R1 +	R2	R3	R4
(Observe Jully developed V 138A	R1 139 B	R2	R3	R4
LEAF PUBESCENCE I 1 = Absent; 2 = Sp	DENSITY: arse; 3 = Medium; 4 =	Thick; 5 = Heavy		
v 3	R1 3	R2	R3	R4
LEAF PUBESCENCE I 1 = None; 2 = Shor	ENGTH: t; 3 = Medium; 4 = La	ong; 5 = Very long		
v 2	R1 2	R2	R3	R4
(Note: Descriptor #19	can be used to describe ti	he type and length of the	glandular trichomes observ	ed.)
E LEAF SILHOUETTE:				
v S	R1 5	R2	R3	R4
D-476 (01-96)				Page 4 of 19

		Weak; 5 = Medium; 7		rong	
	v	R1	R2	R3	R4
	LEAF STIPULES SIZ 1 = Absent; 3 = 5	Æ: (See figure 13) Small; 5 = Medium; 7	= Large		
·	V 2	R1 5	R2	R3	R4
	1 = Narrowly ovat	ET SHAPE: (See figure 3 e; 2 = Medium ovate; 3 Oblong; 8 = Other		Lanceolate; 5 = Elliptic	ai;
	v 3	R1 3	R2	R3	R4
		T TIP SHAPE: (See figu uspidate; 3 = Acuminat		her	
-	v 3	R1 3	R2	R3	R4
*	1 = Cuneate; 2 =		•	te; 6 = Lobed; 7 = Oth	er
	v 5	R1 5	R2	R3	R4
*	TERMINAL LEAFLE	T MARGIN WAVINESS light; 3 = Weak; 4 = I			
	v 3	R1 3	R2	R3	R4
	NUMBER OF PRIMA AVERAGE:	RY LEAFLET PAIRS:	(See figure 11)		
	v 3.5	R1 3.5	R2	R3	R4
RA V	NGE: 3 to 4	1 3 to 4	R2 to	R3 to	R4 to
	PRIMARY LEAFLET 1 = Acute; 2 = Ct	TIP SHAPE: (See figur uspidate; 3 = Acuminat	e 4 & 11) e; 4 = Obtuse; 5 = Ot	her	
	v 3	$\begin{bmatrix} \mathbf{R}1 & 3 \end{bmatrix}$	R2	R3	R4
100	476 (01-96)	······································			Page 5 of 19

V 3	1	PRIMARY LEAFLI 1 = Very Small;		ım; 4 = Large; 5 = Ver	v Large	
1 = Narrowly ovate; 2 = Medium ovate; 3 = Broadly ovate; 4 = Lanceolate; 5 = Elliptical; 6 = Obovate; 7 = Oblong; 8 = Other V 2 R1 2 R2 R3 R4 PRIMARY LEAFLET BASE SHAPE: (See figure 5 & 11) 1 = Cuncate; 2 = Acute; 3 = Oblinse; 4 = Cordate; 5 = Truncate; 6 = Lobed; 7 = Other V 5 R1 5 R2 R3 R4 NUMBER OF SECONDARY AND TERTIARY LEAFLET PAIRS: (See figure 11) AVERAGE: V 3 4 R1 3 0 R2 R3 R4 RANGE: V 2 to 5 R1 2 to 4 R2 to R3 to R4 to 5. INFLORESCENCE CHARACTERISTICS: NUMBER OF INFLORESCENCE / PLANT: AVERAGE: V 9 2 R1 3 5 R2 R3 R4 RANGE: V 9 2 R1 3 5 R2 R3 R4 RANGE: V 9 5 to 12 R1 3 to 5 R2 to R3 to R4 to NUMBER OF FLORETS / INFLORESCENCE: AVERAGE: V 5 4 R1 5 7 R2 R3 R4 RANGE: V 5 4 R1 5 7 R2 R3 R4 RANGE: V 5 4 R1 5 7 R2 R3 R4 RANGE: V 5 4 R1 5 7 R2 R3 R4 RANGE: COROLLA INNER SURFACE COLOR CHART VALUE: Royal Horticulture Society Color Chart or Munsell Color Chart (Measure predominant color of newly open flower & circle the appropriate color chart)				1	7 / 7	R4
PRIMARY LEAFLET BASE SHAPE: (See figure 5 & 11) 1 = Cuneate; 2 = Acute; 3 = Obtuse; 4 = Cordate; 5 = Truncate; 6 = Lobed; 7 = Other	Pl	1 = Narrowly ov	ate; 2 = Medium ovate		= Lanceolate; 5 = Ellipti	cal;
1 = Cuneate; 2 = Acute; 3 = Obtuse; 4 = Cordate; 5 = Truncate; 6 = Lobed; 7 = Other		v 2	R1 2	R2	R3	R4
NUMBER OF SECONDARY AND TERTIARY LEAFLET PAIRS: (See figure 11) AVERAGE: V 3.4 R1 3.0 R2 R3 R4 RANGE: V 2 to 5 R1 2 to 4 R2 to R3 to R4 to 5. INFLORESCENCE CHARACTERISTICS: NUMBER OF INFLORESCENCE / PLANT: AVERAGE: V 3 to 12 R1 3.5 R2 R3 R4 RANGE: V 3 to 12 R1 3 to 5 R2 to R3 to R4 to NUMBER OF FLORETS / INFLORESCENCE: AVERAGE: V 5.4 R1 5.7 R2 R3 R4 RANGE: V 5.4 R1 5.7 R2 R3 R4 RANGE: V 4 to 7 R1 3 to 7 R2 to R3 to R4 to ** ** ** ** ** ** ** ** **	PI				cate; 6 = Lobed; 7 = Ot	her
AVERAGE: V 3.4 R1 3.0 R2 R3 R4 RANGE: V 2 to 5 R1 2 to 4 R2 to R3 to R4 to 5. INFLORESCENCE CHARACTERISTICS: NUMBER OF INFLORESCENCE / PLANT: AVERAGE: V 9.2 R1 3.5 R2 R3 R4 RANGE: V 8 to 12 R1 3 to 5 R2 to R3 to R4 to NUMBER OF FLORETS / INFLORESCENCE: AVERAGE: V 5.4 R1 5.7 R2 R3 R4 RANGE: V 5.4 R1 5.7 R2 R3 R4 RANGE: V 4 to 7 R1 3 to 7 R2 to R3 to R4 to ** ** ** ** ** ** ** ** **		v 5	R1 5	R2	R3	R4
RANGE: V 2 to 5 R1 2 to 4 R2 to R3 to R4 to 5. INFLORESCENCE CHARACTERISTICS: NUMBER OF INFLORESCENCE / PLANT: AVERAGE: V 9 2 R1 3 5 R2 R3 R4 RANGE: V 8 to 12 R1 3 to 5 R2 to R3 to R4 to NUMBER OF FLORETS / INFLORESCENCE: AVERAGE: V 5 4 R1 5 7 R2 R3 R4 RANGE: V 4 to 7 R1 3 to 7 R2 R3 R4 RANGE: V 4 to 7 R1 3 to 7 R2 to R3 to R4 to ** ** ** ** ** ** ** ** **	N	A	DARY AND TERTIAR	Y LEAFLET PAIRS: (S	ee figure 11)	
V \(\begin{align*} \lambda \to \\ \lambda \\ \\ \lambda \\ \lambd	[v 3.4	R1 3.0	R2	R3	R4
NUMBER OF INFLORESCENCE / PLANT: AVERAGE: V 9 to 12 R1 3 to 5 R2 to R3 R4 RANGE: V 8 to 12 R1 3 to 5 R2 to R3 to R4 to NUMBER OF FLORETS / INFLORESCENCE: AVERAGE: V 5 14 R1 5 7 R2 R3 R4 RANGE: V 14 to 7 R1 3 to 7 R2 to R3 to R4 to ** COROLLA INNER SURFACE COLOR CHART VALUE: Royal Horticulture Society Color Chart (Measure predominant color of newly open flower & circle the appropriate color chart)			R1 2 to 4	R2 to	R3 to	R4 to
AVERAGE: V 9-2 R1 3-5 R2 R3 R4 RANGE: V 8 to 12 R1 3 to 5 R2 to R3 to R4 to NUMBER OF FLORETS / INFLORESCENCE: AVERAGE: V 5-4 R1 5-7 R2 R3 R4 R4 RANGE: V 4 to 7 R1 3 to 7 R2 R3 R4 R4 RANGE: V 4 to 7 R1 3 to 7 R2 R3 R4 R4 RANGE: V 4 to 7 R1 3 to 7 R2 R3 R4 R4 RANGE: V 4 to 7 R1 3 to 7 R2 COROLLA INNER SURFACE COLOR CHART VALUE: Royal Horticulture Society Color Chart or Munsell Color Chart (Measure predominant color of newly open flower & circle the appropriate color chart)	5. IN	FLORESCENCE C	HARACTERISTICS:			
RANGE: V 9 to 12 R1 3 to 5 R2 to R3 to R4 to NUMBER OF FLORETS / INFLORESCENCE: AVERAGE: V 5.4 R1 5.7 R2 R3 R4 RANGE: V 4 to 7 R1 3 to 7 R2 to R3 to R4 to ** COROLLA INNER SURFACE COLOR CHART VALUE: Royal Horticulture Society Color Chart or Munsell Color Chart (Measure predominant color of newly open flower & circle the appropriate color chart)	N		RESCENCE / PLANT:			
NUMBER OF FLORETS / INFLORESCENCE: AVERAGE: V 5.4 R1 5.7 R2 R3 R4 RANGE: V 4 to 7 R1 3 to 7 R2 to R3 to R4 to ** COROLLA INNER SURFACE COLOR CHART VALUE: Royal Horticulture Society Color Chart (Measure predominant color of newly open flower & circle the appropriate color chart)			R1 3.5	R2	R3	R4
NUMBER OF FLORETS / INFLORESCENCE: AVERAGE: V 5,4 R1 5,7 R2 R3 R4 RANGE: V 4 to 7 R1 3 to 7 R2 to R3 to R4 to ** ** ** ** ** ** ** ** **	RAN	GE:				
AVERAGE: V 5.4 R1 5.7 R2 R3 R4 RANGE: V 4 to 7 R1 3 to 7 R2 to R3 to R4 to ** ** COROLLA INNER SURFACE COLOR CHART VALUE: Royal Horticulture Society Color Chart of Munsell Color Chart (Measure predominant color of newly open flower & circle the appropriate color chart)			2 2			
RANGE: V 4 to 7 R1 3 to 7 R2 to R3 to R4 to ** ** ** ** ** ** ** ** **	v		R1 3 to 5	R2 to	R3 to	R4 to
* COROLLA INNER SURFACE COLOR CHART VALUE: Royal Horticulture Society Color Chart or Munsell Color Chart (Measure predominant color of newly open flower & circle the appropriate color chart)		8 to 12 I			R3 to	R4 to
* COROLLA INNER SURFACE COLOR CHART VALUE: Royal Horticulture Society Color Chart or Munsell Color Chart (Measure predominant color of newly open flower & circle the appropriate color chart)		MBER OF FLORE	TS / INFLORESCENCE	£:		
(Measure predominant color of newly open flower & circle the appropriate color chart)	NU RAN	MBER OF FLORE AVERAGE: V 5.4	TS / INFLORESCENCE R1 5.7	R2	R3	R4
V 155B R1 1558 R2 R3 R4	NU RAN	MBER OF FLORE AVERAGE: V 5.4	TS / INFLORESCENCE R1 5.7	R2	R3	R4
	NU RAN V	MBER OF FLORE AVERAGE: V 5.4 IGE: L to 7 IROCALLA INNER SU	R1 3 to 7 URFACE COLOR CHAI	R2 to RT VALUE: Royal Hort	R3 to	R4 to

* COROLLA OUT (Measure predom)				ral Horticulture Society Co opriate color chart)	olor Chart or Munsell Color Chart
v iss&	R1	155B	R2	R3	R4
STD-476 (01-96)					Page 6.5 of 19

	IER SURFACE COLOI 2 = Red-violet; 3 = Bl		color of newly open flower)	
V	R1	R2	R3	R4
	APE: (See figure 6) ate; 2 = Rotate; 3 = 1	Pentagonal; 4 = Semi-st	ellate; 5 = Stellate	
v 4	R1 4	R2	R3	R4
	OCYANIN COLORATI 3 = Weak; 5 = Mediu	ON: um; 7 = Strong; 9 = V	ery strong	
v	R1	R2	R3	R4
			y Color Chart st Munsell appropriate color chart)	Color Chart
v 9A	RI 9A	R2	R3	R4
	E: (See figure 7) ne; 2 = Narrow cone; R1	3 = Pear shape cone;	4 = Loose; 5 = Other	R4
POLLEN PROD 1 = None; 3	UCTION: = Some; 5 = Abunda	nt		
v 3	R1 3	R2	R3	R4
STIGMA SHAP 1 = Capitate	E: (See figure 8) 2 = Clavate; 3 = Bi-	lobed		
v	R1	R2	R3	R4
	R CHART VALUE (R	oyal Horticulture Society	y Color Chart or Munsell	Color Chart
v 133A	R1 138A	R2	R3	R4
BERRY PRODU 1 = None; 3	CTION: (Under field c = Low; 5 = Moderate	onditions) ; 7 = Heavy; 9 = Very	y heavy	
v 3	R1 3	R2	R3	R4

5. TUBER CHARACTER	dsiics:			·
	KIN COLOR: Light Yellow; 3 = Yello ; 10 = Purple; 11 = Da			8 = Red;
V	R1 4	R2	R3	R4
PREDOMINANT SI (Circle the appropriat	KIN COLOR CHART V. te color chart)	ALUE: Royal Horticultu	re Society Color Chart	or Munsell Color Chart
v 1588	R1 1640	R2	R3	R4
SECONDARY SKIN 1 = Absent; 2 =	COLOR: Present, please describe	•		
v	R1	R2	R3	R4
SECONDARY SKIN (Circle the appropriate	COLOR CHART VALU e color)	JE: Royal Horticulture	Society Color Chart or N	Aunsell Color Chart
V NA	R1 NA	R2	R3	R4
	COLOR DISTRIBUTION COLOR DISTRIBUTION COLOR DISTRIBUTION (COLOR DISTRIBUTION COLOR DISTRI		ectacled; 6 = Stippled;	7 = Other
VNA	R1 NA	R2	R3	R4
SKIN TEXTURE: 1 = Smooth; 2 =	Rough (flaky); 3 = Net	ted; 4 = Russetted; 5	= Heavily russetted; 6 =	= Other
v 1	R1 2	R2	R3	R4
* TUBER SHAPE: (Se 1 = Compressed;	ce figure 10) 2 = Round; 3 = Oval;	4 = Oblong; 5 = Long	g; 6 = Other	
v 3	RI 2	R2	R3	R4
TUBER THICKNES 1 = Round; 2 ==	S: Medium thick; 3 = Slig	htly flattened; 4 = Flat	tened; 5 = Other	
v 2	R1 2	R2	R3	R4

TUBER LENGTH (mm):			
AVERAGE:		[
v 72.9 R1 66.5	R2	R3.	R4
RANGE:			
V 52 to 94 R1 45 to 97	R2 to	R3 to	R4 to
STANDARD DEVIATION:		-	
v 10.9 R1 13.2	R2	R3	R4
A VIEW A CIE WIEWCYM OF CARANTE FRA V			
V 8.20 kg R1 8.70 kg	R2	R3	R4
V 3 20 75	[RZ]	KS	R4
TUBER WIDTH (mm):			
AVERAGE:			
v 65.3 R1 68.5	R2	R3	R4
RANGE:	.		
v 50 to 83 R1 54 to 95	R2 to	R3 to	R4 to
STANDARD DEVIATION:			
v - 8.6 R1 10.4	R2	R3	R4
<u> </u>			
v 3.20kg R1 8.70kg		D2	P4
V 8.20kg R1 8.70kg	R2	R3	R4
TUBER THICKNESS (mm):			
AVERAGE:			
v 49.7 R1 53.0	R2	R3	R4
RANGE:			
v 37 to 64 R1 44 to 66	R2 to	R3 to	R4 to
STANDARD DEVIATION:		•	
v 6.4 R1 6.4	R2	R3	R4
AVERAGE WEIGHT OF SAMPLE TAK	en.		· · · · · · · · · · · · · · · · · · ·
V 8.20 kg R1 8.70 kg	R2	R3	R4
TUBER EYE DEPTH:			
1 = Protruding; 2 = Shallow; 3 = Inter	rmediate; 4 = Deep; 5 =	· Very deep	 1
\mathbf{v} 3 \mathbf{R}_1 3	R2	R3	R4

	TUBER LATERAL EYES 1 = Protruding; 2 = Shallow; 3 = Intermediate; 4	= Deep; 5 = V	ery deep	
	v 3 R1 3 R	2	R3	R4
	NUMBER EYE / TUBER:			
	AVERAGE:	-		
	V 13-0 R1 9-0 R	2	R3	R4
	ANGE:			
V	12 to 14 R1 8 to 10 R2	to	R3 to	R4 to
	DISTRIBUTION OF TUBER EYES: 1 = Predominantly apical; 2 = Evenly distributed			
,	v 2 R1 2 R	2	R3	R4
	PROMINENCE OF TUBER EYEBROWS:			
	1 = Not prominent; 2 = Slight prominence; 3 = Mo	edium promineno	ce; 4 = Very prominent;	5 = Other
	V R1 R	2	R3	R4
*	PRIMARY TUBER FLESH COLOR CHART VALU	F. (Dava) Harris	ultura Society Céla Char	t on Mungali Colon Chant
	(Circle the appropriate color chart)	E: (Royal Horuc	unitie Society Color Char	t or Muisen Color Chart
.]	v 1582 R1 158B R2		R3	R4
	SECONDARY TUBER FLESH COLOR: 1 = Absent; 2 = Present, please describe:	P		
- ".		······································		
	V R1 R	2	R3	R4
	SECONDARY TUBER FLESH COLOR CHART VALU (Circle the appropriate color chart)	E: Royal Hortic	culture Society Color Char	t or Munsell Color Chart
	V NA R1 NA R2		R3	R4
- -	NUMBER OF TUBERS / PLANT: 1 = Low (<8); 2 = Medium (8 -15); 3 = High (>1	5)		
	v 2 R1 2 R	2	R3	R4

DISEASES CHARAC	TERISTICS:			
DISEASES REACT	ION: 0 = NOT TESTE 5 = MODERATE	D; 1 = RESISTANT; 3 LY SUSCEPTIBLE; 7=	= MODERATELY RESIS SUSCEPTIBLE; 9=HIGHI	TANT; AY SUSCEPTIBLE
BACTERIAL RIN	G ROT, FOLIAR REAC	TION:		
v 7	R1 0	R2	R3	R4
BACTERIAL RIN	G ROT, TUBER REACT	TON:		
v 5	Ri O	R2	R3	R4
LATE BLIGHT:				
v 7	R1 0	R2	R3	R4
PLRV (LEAF RO	LL):			
V O	R1 0	R2	R3	R4
PVX:				
VO	R1 O	R2	R3	R4
PVY:				<u> </u>
v O	R1 0	R2	R3	R4
OTHER:				
V	R1 0	R2	R3	R4
OTHER:			,	
V	R1 0	R2	R3	R4
PESTS CHARACTER	ISTICS:			
PEST REACTION:			ODERATELY RESISTANT EPTIBLE; 9=HIGHLY SU	
GOLDEN NEMA	rode:			
v 0	R1 O	R2	R3	R4
OTHER:				
v 0	R1 O	R2	R3	R4
GENE TRAITS:				
INSERTION OF G	ENES: e gene(s) introduced or a	YES	NO NO	
		· · · · · · · · · · · · · · · · · · ·		· · · · · · · · · · · · · · · · · · ·
		· · · · · · · · · · · · · · · · · · ·		

		·						Exhibit	C (P	
NAME OF APPLICANT (S) ICULTURE AND.	ARRI-EDOD TE	EMPORARY OR EXPERIMENTAL DESIGNA	TION	VARIETY N	AME					
DA AND THE		WIS75-30	AC GLACIER CHIF							
WISCONS/N ADDRESS (Street and No. or RD No.,				Tether ?	ก็เกียร	win v				
5403 - 1ST	H, P.O. Box 3000	5	PVPO NUMI	BER						
LETHBRIDGE,	ALBERTA 1	CANADA		#20	0	1 (0 (0	4	
REFERENCE VARIETIES: Er	ster the reference variety name i	n the appropriate box.								
Application Variety (V)	Reference Variety 1 (R1)	Reference Variety 2 (R2)	Reference Varie	ty 3 (R3)	Refe	erenc	e Varie	ety 4 (I	₹4)	
W1575-30	SNOWDEN									
PLEASE READ ALL INSTR	UCTIONS CAREFULLY:					<u> </u>				
. MARKET CHARACTERIST	CS:									
*MARKET CLASS: 1 = Yellow-flesh Table 5 = Russet Tablestock		stock 3 = Chip-processing 4 =	Frozen-processing							
V	R1 I	R2 R3	R4							
LIGHT SPROUT CHARACTI	ERISTICS: (See Figure 1)				·					
*LIGHT SPROUT: G 1 = Spherical 2 =		Panad adiadrina — E — Blossau a	diadrical 5 - Oth	ıer						
1 - Ophelica: 2 -	Ovoid 3 = Conica 4 = E	Broad cylindrica 5 = Narrow cy	minuncai 0 - 00							
v 3	R1 F	22 R3	R4							
*LIGHT SPROUT BAS 1 = Absent 2 = Wo	E: PUBESCENCE OF TIP eak 3 = Medium 4 = S	trong 5 = Very Strong								
V 3	R1 2 R	2 R3	R4							
*LIGHT SPROUT BAS 1 = Green 2 = Red-	E: ANTHOCYANIN COLORA violet 3 = Blue-violet 4	TION = Other(describe)		<u>.</u>						
V 2	R1 / R2	2 R3	R4							
*LIGHT SPROUT BASI 1 = Absent 2 = Wes		ANIN COLORATION (IF PRESEN ong 5 = Very Strong	IT)							
V 5	R1 2 R2	R3	R4							
* LIGHT SPROUT TIP: 1 = Closed 2 = Inte	HABIT ermediate 3 = Open									
V 9	R1 2 R2	R3	R4					-		

Exhibit C (Pota #200100043 2. LIGHT SPROUT CHARACTERISTICS: (continued) LIGHT SPROUT TIP: PUBESCENCE 1 = Absent 2 = Weak 3 = Medium 4 = Strong 5 = Very Strong R4 R3 LIGHT SPROUT TIP ANTHOCYANIN COLORATION 1 = Green 2 = Red-violet 3 = Blue-violet 4 = Other(describe) R3 R2 R4 R1 LIGHT SPROUT TIP: INTENSITY OF ANTHOCANIN COLORATION (IF PRESENT) 2 = Weak 1 = Absent 3 = Medium 4 = Strong 5 = Very Strong R4 R1R₂ R3 LIGHT SPROUT ROOT INITIALS: FREQUENCY 1 = Short 2 = Medium 3 = Long R₂ **R3** R4 R13. PLANT CHARACTERISTICS: **GROWTH HABIT: (See Figure 2)** 3 = Erect (>45° with ground) 5 = Semi-erect (30-45° with ground) 7 = Spreading R1R3 R4 TYPE: 1 = Stem (foliage open, stems clearly visible) 2 = Intermediate 3 = Leaf (Foliage closed, stems hardly visible) **R3** R1R2 R4 MATURITY: Days after planting (DAP) at vine senescence


V	R1	R2	R3	R4
---	----	----	----	----

PLANTING DATE:

								<u> </u>	
	7.7		D 1		l Do	10:3	2	1 R4	1
i	V	1	KI	!!!	KZ	10.	'	1.	
- 1			L						

*REGIONAL AREA:

3 = North East (ME, NY, PA, NJ, MD, MA, RI,) 2 = North Central (ND, WI, MI, MN, OH) 1 = Pacific North West (WA, OR, ID, CO, CA) 4 = Mid-Atlantic Erect (VI, NC, SC, South NJ, FL) 5 = South (LA, TX, AZ, NE) 6 = Canada 10 = Brazil 11 = Other 7 = Europe 8 = England 9 = Latin America





1 = Very Early (<100 DAP) 2 = Early (100-110 DAP) 3 = Mid-season (111-120 DAP) 4 = Late (121-130 DAP) 5 = Very Late (>130 DAP).

V	R1	R2	R3	R4
	J []		<u> </u>	

VITY (wt. air /wt. = 1.060-1.069; 3 R1	= 1.070-1.079; 4	3 = 1.080-1.089; s	5 > 1.090 R3	R4
	5 1	છ	R3	R4
OALKALOID CON				
	TENT (mg. / 106	g. fresh tuber)	,	
R1 14	5 [1	22	R3	R4
ing, french fry proc				
TIEICATION.			· · · · · · · · · · · · · · · · · · ·	
	ding protocol.			
	TAR A CORPUS COM	~~		
			41 3: 3 -4:	
-	nat would be user	ai in distinguisini	g the canologie variety.	
			•	
, ;	rubescan	CE - WEAK		
		E - VIC BUE-SE	CHETAI - LIGHT	E RED VIOLET
lacier chil		ac - Pighent		TAJOIN JULB
	Sing, french fry proc rotocol. NTIFICATION: al traits of the candita and the correspon N A DMMENTS AND CH itional descriptors the candital can	Sing, french fry processing, baking, be rotocol. NTIFICATION: al traits of the candidate variety that ta and the corresponding protocol. NA A DIMMENTS AND CHARACTERISTIC itional descriptors that would be usef ROUT AGER CHIP: SHAPE — CHER CHIP: PUSESCENCE CHER CHIP: PUSESCENCE	Sing, french fry processing, baking, boiling, after-cooking rotocol. NTIFICATION: al traits of the candidate variety that aid in its identificate and the corresponding protocol. NA A DIMMENTS AND CHARACTERISTICS: itional descriptors that would be useful in distinguishing the continuous that we would be useful in distinguishing the continuous that we would be useful in distinguishing the continuous that we would be useful in distinguishing the continuous that we would be useful in distinguishing the continuous that we would be useful in distinguishing the continuous that we would be useful in distinguishing the continuous that we would be useful in distinguishing the continuous that we would be useful in distinuous the continuous that we would be useful in distinuous the continuous that we would be useful in distinuous the continuous that we would be useful in distinuous the continuous that we would be useful in distinuous the continuous that we would be useful in distinuous the continuous that we wou	NTIFICATION: al traits of the candidate variety that aid in its identification (e.g. protein or DN ta and the corresponding protocol. N Q DMMENTS AND CHARACTERISTICS: itional descriptors that would be useful in distinguishing the candidate variety. ROUT AGEL CHIP: SHAPE - CONICAL EN : SHAPE - SPRENICAL CHER CHIP: PUBESCENICE - MEDIUM

Figure 1: Growth Habit








Figure 2: Leaf Silhouette







Figure 3: Terminal Leaflet Shape / Primary Leaflet Shape



1=Narrowly
Ovate



2=Medium Ovate



3=Broadly Ovate



4=Lanceolate



5=Elliptical



6=Obovate



7=Oblong

Figure 4: Terminal Leaflet Shape of Tip / Primary Leaflet Shape of Tip




Figure 5: Terminal Leaflet Shape of Base / Primary Leaflet Shape of Base





Figure 6: Corolla Shape





stellate |>b




semi-stellate | = b



pentagonai I < b



rotate



very rotate

Figure 7: Anther Shape



1=Broad cone



2=Narrow cone



3=Pear shape cone



4=Loose

Figure 8: Stigma Shape







2=Clavate



3=Bilobed

Figure 9: Distribution of Secondary Tuber Color



1 = Eyes



2=Eyebrows



3=Splashed



4=Scattered



5=Spectacled



6=Stippled

Figure 10: Tuber Shape




Figure 11: Leaf Dissection




Figure: 12 Stem Wings





Figure 13: Leaf Stipules:



2,00100043

REPRODUCE LOCALLY. Include form number and edition date on all	all reproductions. FORM APPROVED - OMB No. 0581-0055			
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). The information is held confidential until the certificate is issued (7 U.S.C. 2426).			
NAME OF APPLICANT(S) Agriculture and Agri-Food Canada and Wisconsin	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME		
Alumni Research Foundation	WIS 75-30	AC Glacier Chip		
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code)	6. FAX (Include area code)		
c/o The Director, Agriculture and Agri- Food Canada, Research Centre, 5403-1st Ave South, Lethbridge, Alberta, Canada	403-327-4561 7. PVPO NUMBER	403-382-3156		
T1J 4B1	200/00043			
8. Does the applicant own all rights to the variety? Mark an "X" in the	appropriate block. If no, please exp l	ain. YES NO		
9. Is the applicant (individual or company) a U.S. national or a U.S. ba	ased company? If no, give name of	country. YES X NO		
Canada 10. Is the applicant the original owner? YES	, NO If no, please answer one	of the following:		
To the applicant the original officer.	X 110, please allswer bill	joi the following.		
b. If the original rights to variety were owned by a company(ies),	is (are) the original owner(s) a U.S. b	ased company?		
11. Additional explanation on ownership (If needed, use the reverse for	or extra space):			
The original cross (1975) was made by The and early generation selection conducted i cross were provided to Agriculture and Agrand the final selection was make by the Le agreement between Agriculture and Agri-Foo Foundation was signed on November 23, 1998 exclusive right to pursue protection of the	n Rhinelander, Wiscons i-Food Canada (Lethbri thbridge Research Cent d Canada and The Wisco granting Agriculture	in. Selected clones of the dge Research Centre) in 198 re. An inter-institutional nsin Alumini Research		
PLEASE NOTE:		·		
Plant variety protection can only be afforded to the owners (not license	ees) who meet the following criteria:			
If the rights to the variety are owned by the original breeder, that penational of a country which affords similar protection to nationals of	rson must be a U.S. national, national the U.S. for the same genus and spec	of a UPOV member country, or cies.		
If the rights to the variety are owned by the company which employed nationals of a UPOV member country, or owned by nationals of a co- genus and species.	ed the original breeder(s), the compar ountry which affords similar protection	ny must be U.S. based, owned by to nationals of the U.S. for the same		
3. If the applicant is an owner who is not the original owner, both the o	riginal owner and the applicant must	meet one of the above criteria.		
The original breeder/owner may be the individual or company who dis-	and and the final burneding. One Occasion	41/aV2) at the Dient Veriety Dretection		

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid CMB control number. The valid CMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 6 minutes per response, including the time for reviewing the instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.

Act for definitions.

REPRODUCE LOCALLY, Include form number and date on all reproductions.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid GMB control number. The ref
CMB control number to this information collection is of\$1-0055. The time required to complete this information collection of information unless per response, including the time for reviewing instructions,
searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the besis of race, color, national origin, gender, religion, age, disability, sexual orientation, market or tamily status politicis, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require atternative means for communication of program information (Braille, large print, eudiolage, etc.) should contact USDA's TARGET Center at 202-720-2000 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or Call 202-720-5964 (voice and TE USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY ANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

EXHIBIT F DECLARATION REGARDING DEPOSIT

NAME OF OWNER (S)

Agriculture and Agri-Food Canada and

NAME OF OWNER REPRESENTATIVE (S) 18-2-07

AGRICULTURE AND MERI-FOOD

CAN ADA (LETHBAIDGE)

ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country)

5403 - IST AVENUE SOUTH, P.O.

Wisconsin Alumni Research Foundation box 3000 , LETHER 1065 , ALBERTA CANA DA

ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country)

5403-1ST AVENUE SOUTH P.O. BOX 3000 (LETHBAIDGE ALBERTA , CANADA

TEMPORARY OR EXPERIMENTAL DESIGNATION

AC GLACIER Per President

#2⁰00100043

I do hereby declare that during the life of the certificate a viable sample of propagating material of the subject variety will be deposited, and replenished as needed periodically, in a public repository in the United States in accordance with the regulations established by the Plant Variety Protection Office.

Mar 13/07